

INTERNATIONAL **BEST PRACTICE RECOMMENDATIONS**

PREVENTION AND
MANAGEMENT OF
**MOISTURE-ASSOCIATED
SKIN DAMAGE
(MASD)**

RECOMMENDATIONS FROM AN EXPERT WORKING GROUP



Wounds
INTERNATIONAL

PUBLISHED BY:
Wounds International
108 Cannon Street
London EC4N 6EU, UK

Tel: + 44 (0)20 3735 8244
info@woundsinternational.com
www.woundsinternational.com

© Wounds International, 2020



The meeting and this document have been supported by 3M.

3M Science.
Applied to Life.

The views in this document do not necessarily reflect those of the sponsor.

How to cite this document:

Fletcher J, Beeckman D,
Boyles A et al (2020)
International Best Practice
Recommendations: Prevention
and management of moisture-
associated skin damage
(MASD). Wounds International.
Available online at www.
woundsinternational.com

EXPERT PANEL

Jacqui Fletcher (Co-chair), Independent Nurse Consultant, UK

Dimitri Beeckman (Co-chair), PhD, RN, FEANS, Professor at Ghent University, Monash University, University of Southern Denmark, Örebro University, University of Surrey, and Royal College of Surgeons in Ireland

Anna Boyles, Lead Clinical Nurse Specialist, Stoma Care, King's College Hospital NHS Foundation Trust, London, UK

Sian Fumarola, Clinical Specialist, NHS Supply Chain; Senior Clinical Nurse Specialist, Tissue Viability and Continence, University Hospitals of North Midlands NHS Trust, Staffordshire, UK

Jan Kottner, Professor, Charité-Universitätsmedizin Berlin, Department of Dermatology and Allergy, Clinical Research Center for Hair and Skin Science, Berlin, Germany; Department of Public Health, Faculty of Medicine and Health Sciences, Ghent University, Belgium

Laurie McNichol, MSN, RN, CNS, GNP, CWOCN, CWON-AP, FAAN, Clinical Nurse Specialist, WOC Nurse, Cone Health, Greensboro, North Carolina, USA

Zena Moore, School of Nursing and Midwifery, Royal College of Surgeons in Ireland (RCSI), University of Medicine and Health Sciences, Dublin, Ireland; Skin Wounds and Trauma (SWaT) Research Center, Royal College of Surgeons in Ireland (RCSI), University of Medicine and Health Sciences, Dublin, Ireland; Faculty of Medicine, Nursing and Health Sciences, Monash University, Melbourne, VIC, Australia; Department of Public Health, Faculty of Medicine and Health Sciences, Ghent University, Belgium; Lida Institute, Shanghai, China; Cardiff University, UK

Nikki Sarkar, Practice Development Nurse in Adult Critical Care, Nottingham University Hospitals NHS Trust, UK

Madeleine Stenius, Training & Education Coordinator/Assistant Nurse, Aleris Rehab Station, Stockholm, Sweden

David Voegeli, Professor of Nursing, Faculty of Health & Wellbeing, University of Winchester, UK

FOREWORD

Moisture-associated skin damage (MASD) represents a significant problem and can have a negative effect on patient wellbeing and quality of life.

A group of international experts met online via Zoom in June 2020 to discuss the key issues and knowledge gaps in MASD, and to formulate Best Practice Statements to guide the prevention and management of MASD.

For the purposes of this document, the wider term of MASD has been subdivided into key areas, with specific Best Practice Statements around prevention and management for each area.

These are:

- Incontinence-associated dermatitis (IAD)
- Peristomal dermatitis
- Intertriginous dermatitis (intertrigo)
- Periwound maceration.

During the meeting, we agreed to keep the umbrella term 'MASD', in order not to overload busy clinicians with differing terminology. However, it is important to note that 'moisture' is not the only relevant factor in these areas of skin damage, as explained in the document.

The aim of this document is to provide clinicians with guidance for best practice in these areas of MASD, thereby contributing to improving patient outcomes and reducing the incidence of these and related skin conditions, through skin-focused prevention and management plans.

Jacqui Fletcher and Dimitri Beeckman (Co-chairs)

Overview: Key issues in MASD

MASD is a complex and increasingly commonly recognized condition. Overexposure of the skin to bodily fluids can compromise its integrity and barrier function, making it more permeable and susceptible to damage (Gray et al, 2001; Woo et al, 2017). Individuals with MASD experience persistent symptoms that affect quality of life, including pain, burning and pruritis (Gray et al, 2011; Woo et al, 2017).


MASD is classified as an irritant-contact dermatitis; see Table 1 (WHO, 2020). Common irritants can include urine, stool, perspiration, saliva, intestinal liquids from stomas and exudate from wounds. As such, MASD is an umbrella term and forms of MASD may be subdivided into four types (see Figure 1):

- IAD
- Peristomal dermatitis (relating to colostomy, ileostomy/ ileal conduit, urostomy, suprapubic catheter, or tracheostomy)
- Intertriginous dermatitis (intertrigo: where two skin areas may touch or rub together)
- Periwound maceration.

Table 1. Types of irritant contact dermatitis according to WHO ICD-10 coding

L24.A Irritant contact dermatitis due to friction or contact with body fluids
L24.A0 Irritant contact dermatitis due to friction or contact with body fluids, unspecified
L24.A1 Irritant contact dermatitis due to saliva
L24.A2 Irritant contact dermatitis due to fecal, urinary or dual incontinence
L24.A9 Irritant contact dermatitis due friction or contact with other specified body fluids
L24.B Irritant contact dermatitis related to stoma or fistula
L24.B0 Irritant contact dermatitis related to unspecified stoma or fistula
L24.B1 Irritant contact dermatitis related to digestive stoma or fistula
L24.B2 Irritant contact dermatitis related to respiratory stoma or fistula
L24.B3 Irritant contact dermatitis related to fecal or urinary stoma or fistula

FIGURE 1 | Types of MASD (Gray et al, 2011)



The development of MASD involves more than bodily fluids alone. Rather, skin damage is attributable to multiple factors, including chemical irritants within the moisture source (e.g. proteases and lipases in feces, drug metabolites), its pH, associated microorganisms on the skin surface (e.g. commensal skin flora), and mechanical factors such as friction (Gray et al, 2011).



Risk assessment and prevention strategies are of key importance in MASD. Interventions can be taken to protect the skin and prevent MASD, including the use of skin protection products, such as barrier creams, liquid polymers, and cyanoacrylates to create a protective layer on the skin surface that simultaneously maintains hydration levels while blocking external moisture and irritants (Gray et al, 2011; McNichol et al, 2018).

Emerging evidence now highlights the links between MASD and other skin conditions such as cutaneous infection and pressure ulcers (Jones et al, 2008; Beeckman et al, 2014). Adopting a holistic, integrated approach, focused on prevention strategies and the importance of skin integrity, can have overall beneficial results and help to break down barriers to effective care in practice (Beeckman et al, 2020).

Clinicians must be vigilant, both in maintaining optimal skin conditions and in diagnosing and treating early stages of MASD to prevent progression and skin breakdown (Gray et al, 2011).

Incontinence-associated dermatitis (IAD)

The term incontinence-associated dermatitis (IAD) describes the skin damage associated with exposure to urine, stool or a combination of these in adults (ICD 10 **L24.A2**). In babies or small children, it is also known as diaper or nappy rash (ICD 10 **L24.A2**), among other terms. The term IAD is preferred, since it distinguishes skin problems arising directly from contact with urine and/or feces due to incontinence from other conditions. Further, IAD acknowledges that the condition may affect more than the perineal area and people of any age (Beeckman et al, 2015).

IAD can have a significantly detrimental effect on patients' quality of life, causing considerable discomfort and in some cases both physical and mental distress (Van den Bussche et al, 2018). From a clinician point of view, it can be difficult, time-consuming and expensive to manage in practice (Doughty et al, 2012; Beeckman et al, 2014).

When identifying risk, consider causal, indirect and contextual factors

Best Practice Statement

Identifying and managing risk

Identifying those at risk, and implementing prevention care, are key in IAD (Beeckman et al, 2015). When considering risk factors and assessing patient risk, it is important to consider causal, indirect and contextual factors (see Table 2). A causal, or direct, risk factor means that one of these risk factors must be present for the diagnosis of IAD to be made.

Table 2. Examples of causal and indirect risk factors for IAD (adapted from Beeckman et al, 2015)

Causal	<ul style="list-style-type: none"> ■ Type of incontinence <ul style="list-style-type: none"> - Urine, feces or both - Solid or liquid (liquid stool poses greater risk)
Indirect	<ul style="list-style-type: none"> ■ Use/non-use of diapers ■ Exposure time ■ Frequency and volume ■ Some foods/drugs in urine or stool ■ Mechanical force (e.g. based on positioning) ■ Poor skin condition ■ Type and frequency of washing ■ Use of occlusive containment products/cleansing ■ Compromised mobility ■ Diabetes ■ Increased age ■ Psychosocial factors ■ Diminished cognitive awareness ■ Inadequate personal hygiene ■ Medication (e.g. antibiotics, immunosuppressants) ■ Malnutrition or infrequent/inadequate intake of food or fluids ■ Smoking ■ Critical illness ■ Fever ■ Low oxygen saturation



Identifying patients at increased risk

In assessing potential risk factors, an observational study in critically ill patients with fecal incontinence (Van Damme et al, 2018) found that factors including liquid stool, diabetes, advanced age, smoking, non-use of diapers, fever, and low oxygen saturation were independently associated with IAD. However, accurate risk assessment and classification tools are required (Beeckman et al, 2015).

In the critical care setting, data suggest that fecal incontinence may be an underestimated problem (with surveyed prevalence between 9 to 37% of patients), which is associated with a high use of nursing time (Bayón Garcia et al, 2011). Patients with fecal incontinence in this setting commonly had compromised skin integrity: perineal dermatitis, moisture lesions or sacral pressure ulcers. However, staff reported moderately low awareness levels of the clinical challenges involved in management and use of fecal management systems was low.

Among staff where awareness was higher, key reported benefits of fecal management systems included: reduced risk of cross-contamination and infection, reduced risk of skin breakdown, and improved patient comfort and dignity (Bayón Garcia et al, 2011).

Managing continence

As a priority, wherever possible, the cause of incontinence should be identified and eliminated, and treatment options examined if possible – although this may be due to a range of factors including health conditions and mobility issues (Wishin et al, 2008; Beeckman et al, 2020). This should include evaluation of bladder and kidney function regarding urinary incontinence, and that of the intestine and colon in the case of fecal incontinence (Beele et al, 2017).

If continence enhancement is not possible, suitable incontinence products should be used and non-invasive behavioural interventions implemented (Beeckman et al, 2018). Behavioural interventions may include nutritional and fluid management, mobility enhancement, and use of different toileting techniques (Wishin et al, 2008; Beeckman et al, 2020).

While IAD does not only affect elderly people, evidence from studies involving elderly nursing home residents suggests that structured toileting and exercise interventions can improve incontinence (Bates-Jensen et al, 2003; Beeckman et al, 2020). The type and frequency of incontinence should be re-assessed on regular basis, in order to tailor incontinence management strategies to the individual and assess the risk of skin-related damage (Beeckman et al, 2018).

Wherever possible, indirect risk factors should be mitigated. This may involve addressing comorbidities or any psychosocial issues that can be optimized (Beeckman et al, 2020).

Skin cleansing

It is recommended to limit exposure to cleansing agents and to use soap substitutes and leave-on products with physical protective properties. Mild, low-irritating surfactants and low pH-cleansers should be used, in conjunction with lukewarm water and soft cloths (Lichterfeld-Kottner et al, 2020).

It is vital to recognize patients at increased risk of developing IAD (e.g. liquid stool or diarrhea) and implement appropriate prevention strategies

Best Practice Statement

The cause of incontinence should be identified and managed

Best Practice Statement

Use a cleanser with a mild surfactant and consider cleansing techniques; soap and water should be avoided to maintain the skin's normal pH

Best Practice Statement

For patients with IAD, use a skin protectant that can alleviate pain or improve comfort; for patients at risk for IAD, use a skin protectant to repel moisture and irritants

Best Practice Statement

The skin of patients who are incontinent should be cleansed at least once daily and after each episode of fecal incontinence (Beeckman et al, 2015).

Skin protection

It is important to note the difference between moisturisers and moisture barriers: moisturisers are intended to hydrate the skin, whereas moisture barriers should repel moisture and irritants (All Wales Tissue Viability Forum and All Wales Continence Forum, 2014). Moisture barriers can be delivered as polymeric films, creams, ointments or pastes. However, limited evidence is available comparing the efficacy of moisture barriers (Beeckman et al, 2016). Polymer-based barriers have the advantage of being both waterproof and breathable; in contrast to pastes and ointments, which are occlusive.

The performance of the principal ingredients will vary according to the overall formulation and usage. All products should be used according to manufacturers' instructions.

In patients with IAD and mobility issues, there may be a need to use dressings on fragile areas or wounds that are in contact with surfaces while sitting or lying down (e.g. wheelchair cushions or beds), due to friction and shear in connection with transfers and changes of position. Acrylate terpolymer and elastomeric barrier film products can help to protect against friction, which might be a consideration for use in these patients.

Classification and documentation

Accurate diagnosis should differentiate IAD from other potential conditions or causes. It should be noted that there is an association between IAD, its most important etiological factors (incontinence and moisture), and PUs (Beeckman et al, 2014). Studies on this have noted that there may be confusion, with IAD being classified as a PU (Beeckman et al, 2014). See Table 3 for further information on differentiating IAD and PUs.

Various classification tools are in use for IAD, and the lack of consistency in language and terminology can cause issues with classification. The Ghent Global IAD Classification Tool (GLOBIAD) may be used to aid consistency (Beeckman et al, 2018). See Table 4 for classification using the GLOBIAD tool.

Table 3. Differentiation between IAD and pressure ulcer (adapted from Back et al, 2011 and Beeckman et al, 2011; published by Wounds International, 2015)

Parameter	IAD	Pressure ulcer
History	Urinary and/or fecal incontinence	Exposure to pressure/shear
Symptoms	Pain, burning, itching, tingling	Pain
Location	Affects perineum, perigenital, peristomal area; buttocks; gluteal fold; medial and posterior aspects of upper thighs; lower back; may extend over bony prominence	Usually over bony prominence or associated with location of a medical device
Shape/edges	Affected area is diffuse with poorly defined edges/ may be blotchy	Distinct edges or margins
Presentation/ depth	Intact skin with erythema (blanchable/non-blanchable), with/without superficial/ partial-thickness skin loss	1. Presentation varies from intact skin with non-blanchable erythema to full-thickness skin loss 2. Base of wound may contain non-viable tissue
Other	Secondary superficial skin infection (e.g. candidiasis) may be present	Secondary soft tissue infection may be present




Table 4. Ghent Global IAD Categorization Tool (Beeckman et al, 2018)

Category 1: Persistent redness	Category 2: Skin loss
<p>1A - Persistent redness without clinical signs of infection</p> <p>Critical criterion</p> <ul style="list-style-type: none"> Persistent redness A variety of tones of redness may be present. Patients with darker skin tones, the skin may be paler or darker than normal, or purple in color. <p>Additional criteria</p> <ul style="list-style-type: none"> Marked areas or discoloration from a previous (healed) skin defect Shiny appearance of the skin Macerated skin Intact vesicles and/or bullae Skin may feel tense or swollen at palpation Burning, tingling, itching or pain 	<p>2A - Skin loss without clinical signs of infection</p> <p>Critical criterion</p> <ul style="list-style-type: none"> Skin loss Skin loss may present as skin erosion (may result from damaged/eroded vesicles or bullae), denudation or excoriation. The skin damage pattern may be diffuse. <p>Additional criteria</p> <ul style="list-style-type: none"> Persistent redness A variety of tones of redness may be present. Patients with darker skin tones, the skin may be paler or darker than normal, or purple in color Marked areas or discoloration from a previous (healed) skin defect Shiny appearance of the skin Macerated skin Intact vesicles and/or bullae Skin may feel tense or swollen at palpation Burning, tingling, itching or pain
<p>1B - Persistent redness with clinical signs of infection</p> <p>Critical criteria</p> <ul style="list-style-type: none"> Persistent redness A variety of tones of redness may be present. Patients with darker skin tones, the skin may be paler or darker than normal, or purple in color. Signs of infection Such as white scaling of the skin (suggesting a fungal infection) or satellite lesions (pustules surrounding the lesion, suggesting a Candida albicans fungal infection). <p>Additional criteria</p> <ul style="list-style-type: none"> Marked areas or discoloration from a previous (healed) skin defect Shiny appearance of the skin Macerated skin Intact vesicles and/or bullae The skin may feel tense or swollen at palpation Burning, tingling, itching or pain 	<p>2B - Skin loss with clinical signs of infection</p> <p>Critical criteria</p> <ul style="list-style-type: none"> Skin loss Skin loss may present as skin erosion (may result from damaged/eroded vesicles or bullae), denudation or excoriation. The skin damage pattern may be diffuse. Signs of infection Such as white scaling of the skin (suggesting a fungal infection) or satellite lesions (pustules surrounding the lesion, suggesting a Candida albicans fungal infection), slough visible in the wound bed (yellow/brown/greyish), green appearance within the wound bed (suggesting a bacterial infection with Pseudomonas aeruginosa), excessive exudate levels, purulent exudate (pus) or a shiny appearance of the wound bed. <p>Additional criteria</p> <ul style="list-style-type: none"> Persistent redness A variety of tones of redness may be present. Patients with darker skin tones, the skin may be paler or darker than normal, or purple in color Marked areas or discoloration from a previous (healed) skin defect Shiny appearance of the skin Macerated skin Intact vesicles and/or bullae Skin may feel tense or swollen at palpation Burning, tingling, itching or pain

Peristomal dermatitis

There is a need to standardize and promote better understanding of language and terminology relating to peristomal dermatitis. The term 'stoma' refers to any surgically created opening made into a hollow organ, especially one on the surface of the body leading to the gut or trachea. An abdominal stoma is formed by exteriorizing part of the bowel onto the abdominal wall to allow waste to be diverted into a pouch worn on the abdomen. Stomas can arise from any part of the GI tract, and urinary stomas are formed by connecting the ureters to a section of ileum that is resected from GI continuity and used as a conduit; a tracheostomy is a stoma in the true sense of the word but is created in order to maintain the airway.

Peristomal dermatitis refers to skin damage where there is a clear interaction between the skin and the stoma effluent/fluids/secretion/output. Peristomal dermatitis results in inflammation or erosion of the skin due to moisture from fecal, urinary, and chemical irritants beginning at the mucocutaneous junction, which can then spread outwards to affect the surrounding skin. More than 50% of individuals with ostomies experience leakage (Woo et al, 2017). A further study identified that approximately one-third of patients with a stoma who were followed up over a 5-year study period had evidence of skin complications within 90 days of surgery (Taneja et al, 2017). Patients living with an ileostomy are more likely to experience peristomal MASD than were patients living with a colostomy (Colwell et al, 2017; Nagano, 2019).

Risk factors

The following factors increase the risk of developing peristomal dermatitis (Hoeflok et al, 2017):

- Abdominal anatomy: creasing of the skin when changing positions (standing, sitting, supine)
- Location of stoma (e.g. in the GI tract)
- Stoma construction, including degree of protrusion and position of the lumen on the abdomen
- Incorrect pouch, changing technique and/or wear time
- Increased perspiration or exposure to external moisture, which may disrupt the ability of the stoma base plate to fix to the skin, allowing effluent to come into contact with skin (e.g. showering, swimming)
- Incorrect values for how much different types of stoma should be spouted for effective management.

Skin protection and management

The pouching system should be regularly re-evaluated to ensure proper fitting, with the skin barrier suited to the type of output. For abdominal stomas, the first line of protection should be to ensure skin is clean and dry prior to application of the pouch and the focus on ensuring a good fit and seal (i.e. sizing the aperture and fitting with body contours).

When skin damage/dermatitis has occurred, topical products (e.g. skin barrier powders, pastes, rings) may be used to absorb moisture, provide an additional physical barrier, reduce existing irritation, and allow for proper adhesion of the solid hydrocolloid skin adhesive.

Skin protection should be considered in patients prone to skin damage and those who are at risk of developing MARS (medical adhesive-related skin injury). In patients where the peristomal area is denuded, applying stoma appliances can be very challenging and cause discomfort to the patient. Therefore, advanced polymer-based protectants should be applied to protect skin to allow healing and alleviate discomfort. Acrylate terpolymer-based barrier films act as a sacrificial substrate between the skin and the hydrocolloid skin adhesive, and can be considered to prevent MARS.

Skin protection should be considered in patients prone to skin damage and those who have risk of developing MARS

Best Practice Statement



Peri-tube skin protection may be required, for example in the care of individuals with a tracheostomy, gastrostomy or jejunostomy

Best Practice Statement

Where necessary, take steps to identify any additional underlying cause of peristomal dermatitis

Best Practice Statement

Where necessary, use barrier films and adhesive removers to prevent PMARSI

Best Practice Statement

Describe and document the signs and symptoms that are observed in all peristomal dermatitis

Best Practice Statement

If exudate from a wound is the source of moisture, this should be managed with an appropriate absorbent dressing. It is also important to identify the underlying cause in any deeper wounds, which may be related to diseases (e.g. pyoderma gangrenosum), and to exclude pressure damage (e.g. if a belt has been worn too tightly).

Peristomal adhesive-related damage

Skin damage due to MARSIs can occur in stoma care. The term ‘peristomal medical adhesive-related skin injury’ (PMARSI) has been defined as ‘an alteration in skin integrity with erythema and/or other skin alterations such as skin tears, erosion, bulla, or vesicle that is apparent after removal of an adhesive ostomy pouching system’ (Le Blanc et al, 2019). This definition does not include the 30-minute assessment period usually recommended for MARSIs (McNichol et al, 2013), because many individuals with an ostomy cannot leave their ostomy pouching system off for such a prolonged period of time. Le Blanc et al (2019) also state that skin stripping (defined as removing or tearing of the epidermis with removal of the adhesive faceplate) is a particularly prevalent form of PMARSI. Though evidence is lacking, it has been observed that these injuries are frequently associated with unintentional traumatic removal of adhesive products.

Prevention and management strategies in PMARSI are closely linked, including assessment of the individual’s technique when applying or removing the ostomy skin barrier. Management is based on identification of the type of PMARSI – i.e. epidermal stripping, skin tears and tension injuries (Le Blanc et al, 2019). Acrylate terpolymer barrier films can be used to provide a sacrificial layer between the skin and the adhesive stoma device. Adhesive removal products have also proven to be useful.

Classification and documentation

Although there are several in use - e.g. Ostomy Skin Tool (Martins et al, 2010), Ostomy Algorithm (Beitz et al, 2010), Peristomal Lesion Scale (Menin et al, 2018) - there is currently no standardized classification system for peristomal dermatitis or for PMARSI; some classifications do exist, but these are not currently considered fit for purpose. In order to improve care, standardization is required, therefore documentation of peristomal dermatitis is important.

The purpose of this is to:

- Standardize record-keeping in peristomal skin care
- Guide future care and improve outcomes
- Facilitate incident reporting
- Facilitate research.

Further research is needed into prevalence of peristomal dermatitis and PMARSI, and the effect on the individual, therefore documentation is of key importance in this area (Le Blanc et al, 2019; Yates et al, 2017). Working in consultation with specialized ostomy care nurses (SCN, ET or WOC) may also be necessary if issues are not resolved (Colwell et al, 2011).

Intertriginous dermatitis

Intertriginous dermatitis (also known as intertrigo) is a clinical inflammatory condition that develops in opposing skin surfaces in response to friction, humidity, and reduced air circulation (occlusion) – i.e. inflammation resulting from bodily fluids trapped in skin folds subjected to friction (Metin et al, 2018; Sibbald et al, 2013). In the ICD-10, it is described as ‘irritant contact dermatitis of the skin folds (axillary, submammary, genitocrural, abdominal apron) caused by repetitive shearing forces of skin on skin. Sweat, other body fluids, occlusion and obesity all contribute to its development (L24.A)’ (WHO, 2020).

Intertriginous dermatitis can occur in any area of the body where there are two skin surfaces in close contact with each other, such as the interdigital regions of the feet or hands. However, intertriginous dermatitis is more common in the natural large skin folds of the body such as the axillary, inframammary, umbilical, perianal and inguinal areas (Kalra et al, 2014; Metin et al, 2018).

Intertriginous dermatitis tends to be a neglected area of MASD in the literature. A German population-based study indicated that every sixth aged nursing home resident is affected by intertriginous dermatitis (Gabriel et al, 2019). In the Netherlands, overall prevalence is highest in home care, at approximately 10%, followed by nursing homes at approximately 7% (Kottner et al, 2020). Well-designed clinical trials available to support therapies commonly used to treat or prevent intertriginous dermatitis are lacking (Mistiaen and van Halm-Walters, 2010; Sibbald et al, 2013). Further, despite the plethora of treatments, there is a lack of evidence about their efficacy (Black et al, 2011; Mistiaen and van Halm-Walters, 2010).

Identifying risk factors

A number of risk factors have been considered, including hyperhidrosis, immunodeficiency, diabetes mellitus, immobility, large skin folds, and obesity (Woo et al, 2017); all intertriginous dermatitis risk factors are exacerbated by hot and humid environments (Sibbald et al, 2013). While intertriginous dermatitis can affect a variety of patient groups, it is strongly associated with obesity and skin care dependency (Gabriel et al, 2019; Kottner et al, 2020), and there is a relationship between the degree of obesity and incidence of intertriginous dermatitis (Shareef et al, 2018).

A newer group at high risk of developing intertriginous dermatitis are post-bariatric patients, who have undergone significant weight loss but have been left with large amounts of excess skin and extreme skin folds, which may require skin reduction surgery (Acartuk et al, 2004).

Identification and classification

No formal standardized risk assessment tool for intertriginous dermatitis is currently in use. A prevention program for individuals at risk can include weight loss, a skin-fold hygiene program, and early detection and treatment of recurrences (Sibbald et al, 2013).

Diagnosing the presence of intertriginous dermatitis depends upon the intersection of risk factors being present. Intertriginous dermatitis starts as redness and inflammation on the skin and then is also likely to develop infection. The classic clinical signs of intertriginous dermatitis include mirror-image erythema in the skin folds, accompanied by sensations of itching, stinging and burning.



Keep the skin areas at risk clean and dry and inspect regularly

Best Practice Statement

In the majority of cases, intertriginous dermatitis is associated with fungal or bacterial overgrowth, therefore it is vital to keep the area clean and dry, and to minimize friction. In at-risk patients – particularly those with skin folds (e.g. lymphedema, bariatric patients, those with significant weight loss and excess skin) – it is important to inspect the skin regularly for signs of intertriginous dermatitis developing.

Bacterial load and secondary infection

Patients with secondary infection due to *Candida* often complain of intense itching and the inflamed area has clear margins accompanied by satellite lesions (NICE, 2018), while bacterial infection often presents with 'fiery' red lesions, exudate and odor. If any secondary infection is not resolved, intertriginous dermatitis can progress into more serious soft tissue infections, such as cellulitis, or even lead to sepsis, particularly in diabetic patients with interdigital intertriginous dermatitis in the feet (Black et al, 2011; Kalra et al, 2014).

Emphasize the importance of maintaining good principles of hygiene for individuals at risk

Best Practice Statement

It is important to assess and monitor whether the bacterial load is high. This may be evident in the skin (i.e. intensive or demarcated redness, which may indicate bacterial infection). It is also recommended to monitor odor, as presence of this may indicate a high bacterial load. The importance of following good basic hygiene principles for individuals at risk should be emphasized. If necessary, a swab may be taken to ascertain bacterial load and potential infection (Voegeli, 2020).

Management of intertriginous dermatitis

Proposed principles of management for intertriginous dermatitis (Sibbald et al, 2013) suggest that prevention and treatment of intertriginous dermatitis should maximize the intrinsic moisture barrier function of the skin by focusing on at least one of the following goals:

- Minimize skin-on-skin contact and friction
- Remove irritants from the skin and protect the skin from additional exposure to irritants
- Wick moisture away from affected and at-risk skin (consider wicking products)
- Control or divert the moisture source
- Prevent secondary infection.

Treatments such as drying agents (talc, corn starch), astringents and absorptive materials have been used for intertriginous dermatitis. However, more recent work shows that these agents may not be suitable for use, and may cause further irritation (Janniger et al, 2015). Textiles, such as gauze, fabric or paper towels placed in the skin folds should also be avoided, as although they absorb moisture, they do not allow it to evaporate, thereby retain the moisture and increase the risk of damage (Sibbald et al, 2013; Cunliffe, 2018). Newer moisture-wicking fabrics draw moisture away from the skin towards the outer layer of the fabric to keep the wearer dry and comfortable; these fabrics have an increased surface area, allowing for a greater absorption of water away from the skin and a quicker rate of evaporation. Wicking fabrics are generally used between skin folds and must be properly sited to allow any moisture to evaporate. If no signs of increased bacterial load are present, acrylate terpolymer barrier films can also be used, since they help to reduce the friction between skin folds.

Before applying skin care products, rule out infection

Best Practice Statement

Intertriginous dermatitis can cause significant discomfort and adversely affect patient quality of life. Itch may be a problem, particularly in the presence of fungal infection. Scratching of uncontrolled itch can cause significant skin damage and may result in the transfer of infection to other areas of the body – particularly the eyes. If unmanaged, infection may progress to more severe inflammation, with erosion, oozing, exudation, odor and maceration of the skin, which can significantly decrease quality of life (Metin et al, 2018).

Consider the use of moisture-wicking textiles specifically indicated for skin fold management

Best Practice Statement

While skincare products are not generally recommended, topical antifungal and corticosteroid creams (often combined) may be applied, as can skin barrier films to protect the skin from moisture and reduce friction (Sibbald et al, 2013; Cunliffe, 2018). A structured skincare regimen is recommended in at-risk individuals (Gabriel et al, 2019), and moisture-wicking textiles, specifically developed for skin fold management, are now recommended for the prevention and treatment of intertriginous dermatitis. Further research is required in the possible role of barrier products and films to reduce friction.

It is important that patients are educated about skin fold management and advised to wear supportive garments, in combination with loose-fitting, lightweight clothing of natural fabrics that wick moisture away from the skin and minimize skin-on-skin contact (Sibbald et al, 2013). These natural fabrics should also be used for bed linen, to promote air circulation and absorb moisture vapor. Intertriginous dermatitis of the toes may be prevented by wearing open-toed shoes, although care should be taken to protect the feet of those with diabetes (Janniger et al, 2015).

Periwound maceration



An acrylate terpolymer barrier is effective in preventing peri-wound skin damage in patients with exuding VLU

Best Practice Statement

The condition of the surrounding skin should be included in any formal wound assessment

Best Practice Statement

Any underlying causes for excess exudate should be identified and addressed

Best Practice Statement

While the production of exudate is vital to the wound healing process, if not managed effectively, exudate can cause damage to the periwound (surrounding) skin (WUWHS, 2019). Periwound skin is particularly vulnerable to MASD when drainage volume exceeds the fluid-handling capacity of the dressing. In addition, repetitive application and removal of adhesive tapes and dressings may strip away the periwound stratum corneum, precipitating further skin damage (Colwell et al, 2011; Woo et al, 2017).

The prevalence of periwound maceration is not well documented, but it is acknowledged that its impact is 'substantial', both on individuals and healthcare systems (Woo et al, 2017). One large-scale international survey involving 2,018 patients with chronic wounds found that 25% of respondents experienced pain around the wound, likely from periwound maceration and local inflammatory responses (Price et al, 2008).

Periwound maceration delays overall wound healing and is also correlated with higher pain levels prior to and during dressing changes (Woo et al, 2017).

Use of an acrylate terpolymer barrier film has been found to facilitate the healing of larger wounds without increasing costs; hence, use of an acrylate terpolymer barrier film for periwound skin protection in patients with exuding venous leg ulcers (VLUs) is the preferred treatment strategy (Guest et al, 2012).

Identifying the cause

In order to manage periwound maceration, the cause of excess exudate should be identified. Any management strategy must then address the factors that are contributing to high exudate levels (and potential periwound damage), as well as physically handling the volume of exudate.

Heightened and ongoing inflammation is a potential cause that may be overlooked and should be considered and managed if necessary. In wounds that are not healing, heightened and ongoing inflammation is a likely contributor to increased exudate production. This may also be related to wound infection and/or the presence of biofilm (Schultz et al, 2011; Percival, 2017). It is important to differentiate infection from inflammation. The presence of local edema will also result in a higher output of exudate, where appropriate compression should be used.

Any patient comorbidities, medications or psychosocial factors that may be contributing should also be addressed.

Dressing selection

In exuding wounds that may cause periwound maceration, dressing selection is the mainstay of management. In general, dressings manage fluid by absorbing it and/or allowing it to evaporate from the dressing surface (Wounds UK, 2013).

The dressing should be selected to handle the exudate and draw moisture away from the

Dressing selection should take into account exudate and skin issues and be individualized to the patient and their wound

Best Practice Statement

surrounding skin in order to prevent damage where possible. Consideration should be given to the type of exudate (e.g. viscosity) as well as the volume.

Dressing selection should be individual to the patient, taking into account the management factors required – it may be beneficial to try different dressings to find the correct one for the individual needs of the patient and the clinical scenario (WUWHS, 2019).

In highly exuding lower limb wounds such as VLU, it is important that dressings are used that can handle exudate when used under compression. Consider the effects of gravity when applying the dressing.

Box 8: Properties of the ideal dressing (adapted from WUWHS, 2007; Dowsett, 2011; Vowden et al, 2011)

- Available in a range of shapes and sizes across care settings
- Easy to apply
- Does not require a secondary dressing
- Comfortable/reduces pain/does not cause pain on application
- Conformable
- Prevents leakage and strikethrough
- Absorbs odor
- Stays intact and remains in place during wear
- Suitable for extended wear
- Suitable fluid-handling capacity as per level of exudate
- Retains fluid-handling capacity under compression therapy or when used with an offloading device
- Atraumatic and retains integrity on removal
- Unlikely to cause sensitization or to provoke an allergic reaction
- Cosmetically acceptable and available in a range of colors to match the patient's request
- Does not impede physical activity
- Patient can shower with the dressing in situ
- Incorporates sensors/alerts to feedback on dressing performance, need for change and wound condition
- Inactivates factors that enhance inflammation (i.e. MMPs)
- Cost-effective – considering factors such as the unit cost of dressing versus time taken to change, the potential impact on healing by use of cheaper dressings, how to make the case to procurement

Consider gravitational and pressure aspects of exudate flow when applying the dressing

Best Practice Statement

Dressing application

When using dressings, it is important to consider the issue of exudate and potential periwound maceration in terms of application technique. For instance, consider gravitational and pressure aspects of exudate flow (e.g. at the bottom of a leg wound, or at the sacrum in some patients with mobility issues) when applying the dressing.

Minimize risk of skin trauma through appropriate dressing selection, application/removal and use of skin protectant products

Best Practice Statement

The risk of skin trauma during dressing/device removal should be minimized (WUWHS, 2019). Use of low-adherent or silicone dressings, tapes or securement devices, and application of periwound skin protectant ointments, creams or barrier films may help to protect the skin and reduce the risk of damaging the skin further (Bianchi, 2012). If the periwound skin is inflamed due to the irritant effects of exudate, a topical corticosteroid may be indicated (Woo et al, 2017).



Any cavity dressing products (e.g. rope, ribbon or strip materials) should be confined to the wound and kept away from the surrounding skin; the dressing material should be in contact with the wound bed and should eliminate dead space. However, overpacking of the cavity should be avoided (WUWHS, 2019).

Even if waterproof dressings are used, it is worth considering whether the increased moisture may be due to the patient showering/bathing, or any other lifestyle issues that may impact on the dressing's efficacy.

Skin protection

Skin protection products should be used to protect the periwound skin. Advanced polymer based barriers can be used where exudate levels are very high (e.g. in VLU's), or where dressing wear time may be extended beyond your control (i.e. individuals who do not attend their follow-up dressing appointments regularly). Film-forming barriers may also be considered as part of a treatment regimen where large expanses of adhesive are used and replaced frequently, such as with negative pressure wound therapy (NPWT).

Skin protection products should always be used according to the manufacturer's instructions and should be based on suitability for the patient and their wound. For instance, some skin protection products may interfere with dressing adherence and absorption and should be used only in suitable wounds (WUWHS, 2019).

Skin protection products should be used to prevent damage to the periwound skin

Best Practice Statement

Conclusions

While MASD can be divided into four main categories, there are common contributing factors. Importantly, it is not the moisture that matters, but the chemical composition of the bodily fluids, friction and occlusion. Basic care strategies should focus on risk assessment, prevention of mitigating factors, employing an appropriate and structured care pathway to maintain skin integrity.

In all patients, supported self-care should be encouraged wherever possible, with the individual should be encouraged to engage with their treatment and educated about the importance of skin integrity and protection. Individual capability will need to be considered so that the appropriate level of ability and willingness to be involved can be ascertained. All self-care should be encouraged within a support system and the patient must know how to obtain further advice when needed.

While MASD is increasingly recognized as a significant problem, it is apparent that there are still gaps in knowledge and practice. Further research is needed into the efficacy of products in protecting skin from moisture and irritants, and reducing friction. In many areas of MASD, standardized documentation is not carried out. Standardization and documentation are of key importance for data collection and guiding future care and research.

In many cases, all forms of MASD can have a significant effect on patient wellbeing and quality of life, so it is vital that all care is delivered using best practice and with a patient-centered approach.

References

- Acartuk TO, Wachtman G, Heil B et al (2004) Panniculectomy as an adjuvant to bariatric surgery. *Annals of Plastic Surgery* 53(4): 360-6
- All Wales Tissue Viability Nurse Forum and All Wales Continence Forum (2014) Best Practice Statement on the Prevention and Management of Moisture Lesions Available online at: www.welshwoundnetwork.org/files/5514/0326/4395/All_Wales-Moisture_Lesions_final_final.pdf (accessed 21.07.2020)
- Bates-Jensen BM, Alessi CA, Al Samarrai NR, Schnelle JF (2003) The effects of an exercise and incontinence intervention on skin health outcomes in nursing home residents. *J Am Geriatr Soc* 51(3): 348-55
- Bayón García C, Binks R, De Luca E et al (2011) Prevalence, clinical management and risks associated with acute fecal incontinence in the critical care setting: the FIRST questionnaire survey. *Critical Care* 15: P480
- Beekman D, Campbell J, Campbell K et al (2015) Incontinence-associated dermatitis: Moving prevention forward. *Wounds International*. Available online at: www.woundsinternational.com/resources/details/incontinence-associated-dermatitis-moving-prevention-forward (accessed 6.07.2020)
- Beekman D, Campbell K, Le Blanc K et al (2020) Best practice recommendations for holistic strategies to promote and maintain skin integrity. *Wounds International*. Available online at: <https://www.woundsinternational.com/resources/details/best-practice-recommendations-holistic-strategies-promote-and-maintain-skin-integrity> (accessed 6.07.2020)
- Beekman D, Smet S, Van den Bussche K (2018) Incontinence-associated dermatitis: why do we need a core outcome set for clinical research? *Wounds International* 9(2): 21-5
- Beekman D, Van Damme N, Schoonhoven L et al (2016) Interventions for preventing and treating incontinence-associated dermatitis in adults. *Cochrane Database of Systematic Reviews*. Available online at www.cochranelibrary.com/cdsr/doi/10.1002/14651858.CD011627.pub2/full (accessed 27.08.2020)
- Beekman D, Van den Bussche K, Alves P et al (2018) Towards an international language for incontinence-associated dermatitis (IAD): design and evaluation of psychometric properties of the Ghent Global IAD Categorization Tool (GLOBIAD) in 30 countries. *Br J Dermatol* 178(6):1331-40
- Beekman D, Van Lancker A, Van Hecke A, Verhaeghe S (2014) A systematic review and meta-analysis of incontinence-associated dermatitis, incontinence, and moisture as risk factors for pressure ulcer development. *Research in Nursing & Health* 10.1002/nur.21593
- Beele H, Smet S, Van Damme N, Beekman D (2017) Incontinence-associated dermatitis: pathogenesis, contributing factors, prevention and management options. *Drugs Aging* 35(1): 1-10
- Beitz J, Gerlach M, Ginsburg P et al (2010) Content validation of a standardized algorithm for ostomy care. *Ostomy Wound Manage* 56(10): 22-38
- Bianchi J (2012) Protecting the integrity of the periwound skin. *Wound Essentials* 1: 58-64
- Black JM, Gray M, Bliss DZ et al (2011) MASD part 2: incontinence associated dermatitis and intertriginous dermatitis: a consensus. *J Wound Ost Continence Nurs* 38(4): 359-70
- Colwell JC, Ratcliff CR, Goldberg M et al (2011) MASD part 3: peristomal moisture-associated dermatitis and periwound moisture-associated dermatitis: a consensus. *J Wound Ostomy Continence Nurs* 38(5): 541-53
- Colwell JC, McNichol L, Boarini J (2017) North America Wound, Ostomy, and Continence and Enterostomal Therapy Nurses Current Ostomy Care Practice Related to Peristomal Skin Issues. *J Wound Ostomy Continence Nurs* 44(3): 1-5
- Cunliffe T (2018) Intertrigo: clinical guidance. Primary Care Dermatological Society. Available online at: www.pcids.org.uk/clinical-guidance/intertrigo (accessed 21.07.2020)
- Doughty D, Junkin J, Kurz P et al (2012) Incontinence-associated dermatitis: Consensus statements, evidence-based guidelines for prevention and treatment, current challenges. *J Wound Ostomy Continence Nurs* 39(3): 303-15
- European Pressure Ulcer Advisory Panel, National Pressure Injury Advisory Panel, Pan Pacific Pressure Injury Alliance (2019) Prevention and treatment of pressure ulcers/injuries: Clinical practice guideline. The International Guideline. Emily Haesler (Ed.). EPUAP/NPIAP/PPPIA: p88-9
- Gabriel S, Hahnel E, Blume-Peytavi U, Kottner J (2019) Prevalence and associated factors of intertrigo in aged nursing home residents: a multi-center cross-sectional prevalence study. *BMC Geriatrics* 19(1): 105
- Gray M, Black JM, Baharestani MM et al (2011) Moisture-associated skin damage: overview and pathophysiology. *J Wound Ostomy Continence Nurs* 38(3): 233-41
- Gray M, Kent D, Ermer-Seltun J, McNichol L (2018) Assessment, selection, use, and evaluation of body-worn absorbent products for adults with incontinence: A WOCN Society Consensus Conference. *J Wound Ostomy Continence Nurs* 45(3): 243-64
- Guest JF, Taylor RR, Vowden K, Vowden P (2012) Relative cost-effectiveness of a skin protectant in managing venous leg ulcers in the UK. *J Wound Care* 21(8): 389-94, 396-8
- Hoefloek J, Salvadalena G, Pridham S et al (2017) Use of convexity in ostomy care: Results of an international consensus meeting. *J Wound Ostomy Continence Nurs* 44(1): 55-62
- Janniger CK, Schwartz RA, Szepletowski JC, Reich A (2015) Intertrigo and common secondary skin infections. *Am Fam Physician* 72(5): 833-38
- Jones JE, Robinson J, Barr W, Carlisle C (2008) Impact of exudate and odor from chronic venous leg ulceration. *Nurs Stand* 22(45): 53-8
- Kalra MG, Higgins KE, Kinney BS (2014) Intertrigo and secondary skin infections. *Am Fam Physician* 89(7): 569-73
- Kottner J, Everink I, van Haastregt J (2020) Prevalence of intertrigo and associated factors: A secondary data analysis of four annual multicenter prevalence studies in the Netherlands. *Int J Nurs Stud* 104: 103437
- Le Blanc K, Whiteley I, McNichol L et al (2019) Peristomal medical adhesive-related skin injury: Results of an international consensus meeting. *J Wound Ostomy Continence Nurs* 46(2): 125-36
- Lichterfeld-Kottner A, El Genedy M, Lahmann N et al (2020) Maintaining skin integrity in the aged: A systematic review. *Int J Nurs Stud* 103: 103509
- Martins L, Ayello EA, Claessens I et al (2010) The Ostomy Skin Tool: Tracking peristomal skin changes. *Br J Nurs* 19(15): 960, 932-4
- McNichol LL, Ayello EA, Phearman et al (2018) Incontinence-Associated Dermatitis: State of the Science and Knowledge Translation. *Adv Skin Wound Care* 31(11):502-13
- McNichol L, Lund C, Rosen T, Gray M (2013) Medical adhesives and patient safety: state of the science: consensus statements for the assessment, prevention, and treatment of adhesive-related skin injuries. *J Wound Ostomy Continence Nurs* 40(4):365-80
- Menin G, Barbierato M, Peghetti A et al (2018) Design and validation of a 'Peristomal Lesion Scale' for peristomal skin assessment. *Int Wound J* 16: 960
- Metin A, Dilek N, Bilgili SG (2018) Recurrent candidal intertrigo: challenges and solutions. *Clin Cosmet Investig Dermatol* 11: 175-85
- Mistain P, van Halm-Walters M (2010) Prevention and treatment of intertrigo in large skin folds of adults: a systematic review. *BMC Nursing* 9: 12
- Percival SL (2017) Importance of biofilm formation in surgical infection. *Br J Surg* 104: e85-94
- Price PE, Fagervik-Morton H, Mudge EJ et al (2008) Dressing-related pain in patients with chronic wounds: an international patient perspective. *Int Wound J* 5(2): 159-71
- Schultz GS, Davidson JM, Kirsner RS et al (2011) Dynamic reciprocity in the wound environment. *Wound Rep Reg* 19(2): 134-48
- Shareef RA, Prasad PVS, Kaviarasan PK (2018) Cutaneous markers of obesity. *Int J Res Med Sci* 6(11): 3562-66
- Sibbald RG, Kelley J, Kennedy-Evans KL et al (2013) A practical approach to the prevention and management of intertrigo, or moisture-associated skin damage, due to perspiration: Expert consensus on best practice. *Wound Care Canada*. Available online at: www.woundscanada.ca/docman/public/71-bpr-prevention-management-fall-2013/file (accessed 6.07.2020)
- Taneja C, Netsch D, Rolstad BS et al (2017) Clinical and economic burden of peristomal skin complications in patients with recent ostomies. *J Wound Ostomy Continence Nurs* 44(4): 350-7
- Van Damme N, Clays E, Verhaeghe S et al (2018) Independent risk factors for the development of incontinence-associated dermatitis (Category 2) in critically ill patients with fecal incontinence: a cross-sectional observational study in 48 ICU units. *International Journal of Nursing Studies* 81: 30-9
- Van den Bussche K, Kottner J, Beele H et al (2018) Core outcome domains in incontinence-associated dermatitis research. *J Adv Nurs* 74(7): 1605-17
- Voegeli D (2020) Intertrigo: causes, prevention and management. *Br J Nurs* 29(12): 516-22
- Wishin J, Gallagher TJ, McCann E (2008) Emerging options for the management of fecal incontinence in hospitalized patients. *J Wound Ostomy Continence Nurs* 35(1): 104-10
- Woo KY, Beekman D, Chakravarthy D (2017) Management of moisture-associated skin damage: A scoping review. *Adv Skin Wound Care* 30(11): 494-501
- Woo KY, Coutts PM, Price P et al (2009) A randomized crossover investigation of pain at dressing change comparing 2 foam dressings. *Adv Skin Wound Care* 22: 304-10
- World Union of Wound Healing Societies (2019) Wound exudate: Effective assessment and management. Available online at: www.woundsinternational.com/resources/details/wuwhs-consensus-document-wound-exudate-effective-assessment-and-management (accessed 10.06.2020)
- Wounds UK (2013) Best Practice Statement. Effective exudate management. London: Wounds UK. Available at: www.wounds-uk.com
- Yates S, McNichol L, Heinecke SB, Gray M (2017) Embracing the concept, defining the practice, and changing the outcome: Setting the standard for medical adhesive-related skin injury interventions in WOC Nursing Practice. *J Wound Ostomy Continence Nurs* 44(1): 13-7

